

## In-vitro antimicrobial activity of *Cymbopogon citratus* Stem extracts

**Phang Siao Ze<sup>1</sup>, Chao Xu Yu<sup>1</sup>, Lee Sian Jo<sup>1</sup>, Shivkanya Fuloria<sup>1\*</sup>, Vetriselvan Subramaniyan<sup>2</sup>, Srikumar Chakravarthi<sup>2</sup>, Neeraj Kumar Fuloria<sup>1\*</sup>**

<sup>1</sup>Faculty of Pharmacy, AIMST University, Bedong, Kedah 08100 Malaysia

<sup>2</sup>Faculty of Medicine, Bioscience and Nursing, MAHSA University, Kuala Lumpur 42610, Malaysia

### Correspondence Address\*

Shivkanya Fuloria, [shivkanya\\_fuloria@aimst.edu.my](mailto:shivkanya_fuloria@aimst.edu.my)

Neeraj Kumar Fuloria, [neerajkumar@aimst.edu.my](mailto:neerajkumar@aimst.edu.my)

### Abstract

Essential oils have always been a field of interest attributed to their strong antibacterial properties. The present study was intended to determine the in-vitro antimicrobial and antioxidant activities of *Cymbopogon citratus* stems. The study involved successive extraction of *Cymbopogon citratus* stems by Soxhlet extraction using different solvents such as dichloromethane, acetone, ethanol and methanol in increasing order of polarity to yield different solvents extracts. The extracts were investigated for their antimicrobial potential. Among all extracts, dichloromethane extract exhibited the highest antimicrobial activity followed by ethanolic extract, acetone extract and lastly methanolic extract. The study concludes that dichloromethane extracts of *Cymbopogon citratus* stem possess the highest inhibitory action over the growth of bacterial strains of *Staphylococcus aureus*, *Streptococcus pyogenes*, *Bacillus cereus*, *Escherichia coli*, *Acinetobacter baumannii* and *Neisseria gonorrhoeae*. The high antimicrobial potential of dichloromethane extract of *Cymbopogon citratus* stem may be due to the presence of flavonoidal constituents.

### 1. Introduction

Essential oils have always been a field of interest attributed to their high medicinal value in humans and animals (Ali et al., 2015). The development of microbial resistance and patient tolerance have gradually reduced the efficacy of commercially available antibiotics on one hand and increased the demand for alternative antibiotics (Vetriselvan, Saminathan, Ashok Kumar, and Vinoth, 2019). Medicinal plants are a good source of natural antimicrobial agents attributed to their defence mechanism against microorganisms (Shivkanya, Shilpa, Sangita, and Neeraj, 2009). *Cymbopogon citratus* is a frost-tender clumping perennial grass that is popularly

used as a lemony flavouring in Thai, Vietnamese, Laotian and Cambodian cooking and is widely cultivated in Southeast Asia. *C. citratus* is believed to be a native of Malaysia (Mirghani et al., 2012). In English, the herb is commonly known as lemongrass, citronella grass or fever grass. More recently, the aromatic oils of the plant have been extracted and exported for use in perfumes (Srivastava, Dubey and Mishra, 2013). Lemongrass is common and widespread within its natural range, and also occurs in cultivation. The leaves of *Cymbopogon citratus* have been used in traditional medicine and are often found in herbal supplements and teas. Many effects have been attributed to both their oral consumption and topical use, with modern research supporting many of their alleged benefits. Laboratory studies have shown cytoprotective, antioxidant, and anti-inflammatory properties in vitro as well as antifungal properties (Mirghani, Liyana and Parveen, 2012). A total of 16 chemical constituents accounting for 93.69 % of the oil, were identified in *C. citratus* among which, geranial (27.04 %), neral (19.93 %) and myrcene (27.04 %) were the major constituents. Lemongrass should be harvested at the appropriate level of maturity in order to achieve high quality essential oil and low production cost. Only 13 compounds were present at each of the maturity stage. Among 13 compounds, only 7 compounds ( $\beta$ -myrcene, 3-undecyne, neral, geranial, nerol, geranyl acetate and juniper camphor) had a concentration of greater than 1% (Tajidin, Ahmad, Rosenani, Azimah, and Munirah, 2012). The development of bacterial resistance to conventional antibiotics necessitates the search for new antibacterial agents. Hence the present study was carried out to find out the antibacterial activity of lemongrass oil against the selected pathogenic bacteria.

## 2. Material and methods

The *Cymbopogon citratus* were collected from Laguna Merbok in Sungai Petani, district of Kedah state of Malaysia. The collected plant stems were dried using the oven at 40°C for three consecutive days. The chemical and reagents used in the study were of laboratory grade and were procured from Merck, SD fine, and Sigma Aldrich. For example, dichloromethane, acetone, ethanol, and methanol were used for the extraction of *Cymbopogon citratus*. Muller Hinton Agar and Nutrient Broth were used as media and Gentamicin as standard for antimicrobial test.

### 2.1. Authentication of plant materials (Preparation of herbarium)

The specimen including stems was collected from *Cymbopogon citratus* plant. The specimen was washed with water and rinsed with 70% ethanol in order to remove dirt particles.

The specimen was kept between the newspapers and then pressed under the wooden press. The specimen was kept air-dried for one month. The dried specimen was mounted onto an acid-free paper. The herbarium was labeled with its common name, geographical situation, place and date of collection, and other features of the plant (Calabrese, 2005). The herbarium was then submitted to the Pharmaceutical Chemistry Unit, Faculty of Pharmacy, AIMST University. (Herbarium voucher specimen Accession number AIMST/FOP/15 *Cymbopogon citratus*)

## **2.2. Collection and drying of *Cymbopogon Citratus* stems**

The *Cymbopogon citratus* stems were collected, cleaned with distilled water and dried using oven at 40°C for three days. The dried stems were then crushed into a coarse powder using a blender. Later on, the obtained powder was subjected to different solvent extraction using Soxhlet apparatus (Salome, 2012).

## **2.3. Preparation of *Citrus citratus* stem extracts**

The *Cymbopogon citratus* dried stems powder was subjected to successive extraction with dichloromethane, acetone, ethanol and methanol in an increasing order of polarity. About 15 g *Cymbopogon citratus* dried stem powder was placed into the Soxhlet extraction tube. The Soxhlet apparatus was set up and made ready for extraction. Extra care was needed while setting up the apparatus. A few pieces of porcelain chips were placed to reduce the bumping during extraction. The dried stems powder was successively extracted using different solvents such as dichloromethane, acetone, ethanol and methanol. The reaction was started once the solvent was heated for 25 siphonic cycles. After about 25 siphonic cycles (approximately for 8 hours), a change in the colour of the extraction solvent from yellowish/green to colourless was observed in the Soxhlet extracting tube. Lastly, the solution was collected and evaporated using an evaporating dish to obtain a concentrated pure extract (Azwanida, 2015; Redfern, Kinninmonth, Burdass, and Verran, 2014). The experiment was repeated by using different extraction solvents successively in their order of increasing polarity such as dichloromethane, acetone, ethanol, followed by methanol.

## **2.4. Determination of Antibacterial potential**

### **2.4.1. Preparation of bacterial culture**

Six bacterial strains such as *Staphylococcus aureus*, *Staphylococcus pyogenes*, *Bacillus cereus*: Gram positive bacteria, *Escherichia coli*, *Acinetobacter baumannii*, and *Neisseria gonorrhoeae* were used for the antimicrobial study. The microorganism stock cultures were

maintained and stored at 4°C in the refrigerator. The Nutrient Broth and apparatus used in the transfer of microorganisms were previously autoclaved to ensure the absence of other organisms. Subcultures were prepared by transferring a loopful of colonies of microorganisms from the stock culture into Nutrient Broth and incubated at 37°C for 24 hours in the shaking incubator. Turbidity of the broth indicated the growth of microorganisms (Vetrivelan, 2012).

#### **2.4.2. Agar Well Diffusion method**

The antibacterial potential of the extracts of *Cymbopogon citratus* stems was determined using a well diffusion method. Mueller Hinton agar (MHA) medium was used in this test. Approximately 100 µl bacteria suspension was pipetted on the MHA plate and spread over the agar surface evenly using a sterile L-shaped glass rod. Study involved gentamicin (0.1 mg/ml), dichloromethane extract (1 mg/ml), acetone extract (1 mg/ml), ethanol extract (1 mg/ml), and methanol extract (1 mg/ml). Holes with a diameter of 8-10 mm were punched aseptically with a sterile cork borer on the agar containing bacteria suspension. The volume of approximately 100-200 µl of the gentamicin solution, extracts and phosphate buffer at desired concentration were introduced into the well. The agar plates were incubated at 37°C for 24 hours. The diameters of the zones of inhibition with different extracts were measured in mm. Each test was performed in triplicate and the mean values were recorded (Balouiri, Sadiki, and Ibensouda, 2016; Devillers, Steiman, and Seigle-Murandi, 1989; Jaju, Patil, Durgakar, and Fuloria, 2009).

#### **2.4.3. Agar Disc Diffusion method**

The antibacterial potential of the extracts of *Cymbopogon citratus* stems was determined using the disc diffusion method. Mueller Hinton agar (MHA) medium was used in this test. Approximately 100 µl bacteria suspension was pipetted on the MHA plate and spread over the agar surface evenly using a sterile L-shaped glass rod. Study involved gentamicin (0.1 mg/ml), dichloromethane extract (1 mg/ml), acetone extract (1 mg/ml), ethanol extract (1 mg/ml), and methanol extract (1 mg/ml). Dried and sterilized filter paper discs with 6 mm diameter were suspended into the gentamicin solution and extract solution with the help of forceps. Discs with absorbed extracts were placed on the surface of the agar containing bacteria suspension with sterile forceps. The plates were incubated at 37°C for 24 hours. The diameters of the zones of inhibition by discs with different extracts were measured in mm. Each test was performed in triplicate and the mean values were recorded (Balouiri et al., 2016).

### 3. Results

#### 3.1. Determination of antimicrobial potential

In the present study, the prepared extracts of lemongrass (*Cymbopogon citratus*) stem were evaluated for their antimicrobial potential against various bacterial strains such as *Staphylococcus aureus*, *Streptococcus pyogenes*, *Bacillus cereus*, *Escherichia coli*, *Acinetobacter baumannii* and *Neisseria gonorrhoeae*, using Agar well diffusion and disc diffusion methods for measurement of the zone of inhibition. The prepared extracts of lemongrass (*Cymbopogon citratus*) stem were evaluated for their antimicrobial potential against various bacterial strains using well diffusion and disc diffusion methods. The results so obtained are given in Table 1 and Table 2.

**Table 1:** Zone of inhibition of the lemongrass (*Cymbopogon Citratus*) extracts using well diffusion method

Microorganism	Component	Zone of inhibition (mm)			
		Reading 1	Reading 2	Reading 3	Average value
<i>Staphylococcus aureus</i>	Positive control <i>Gentamicin</i> (1mg/ml)	16	20	14	16.7
	Dichloromethane	-	-	-	-
	Acetone	-	-	-	-
	Ethanol	-	2	-	-
	Methanol	-	-	-	-
	Negative control	-	-	-	-
<i>Streptococcus pyogenes</i>	Positive control <i>Gentamicin</i> (1mg/ml)	18	15	13	15.3
	Dichloromethane	8	-	4	6
	Acetone	-	3	3	3
	Ethanol	-	6	2	4
	Methanol	-	-	3	-
	Negative control	-	-	-	-
<i>Bacillus cereus</i>	Positive control <i>Gentamicin</i> (1mg/ml)	19	16	14	16.3
	Dichloromethane	8	5	10	7.67
	Acetone	-	3	2	2.5
	Ethanol	-	3	-	-
	Methanol	2	1	2	1.67
	Negative control	-	-	-	-

<i>Escherichia coli</i>	Positive control <i>Gentamicin</i> (1mg/ml)	18	18	15	17
	Dichloromethane	19	15	-	17
	Acetone	-	10	2	6
	Ethanol	-	8	2	5
	Methanol	-	2	-	-
	Negative control	-	-	-	-
<i>Acinetobacter baumannii</i>	Positive control <i>Gentamicin</i> (1mg/ml)	14	19	15	16
	Dichloromethane	6	8	-	7
	Acetone	2	8	2	4
	Ethanol	-	3	-	-
	Methanol	2	-	2	2
	Negative control	-	-	-	-
<i>Neisseria gonorrhoeae</i>	Positive control <i>Gentamicin</i> (1mg/ml)	17	18	13	16
	Dichloromethane	14	13	-	13.5
	Acetone	-	9	7	8
	Ethanol	-	10	-	-
	Methanol	-	2	3	2.5
	Negative control	-	-	-	-

**Table 2:** Zone of inhibition of the lemongrass (*Cymbopogon Citratus*) extracts using disc diffusion method

Microorganism	Component	Zone of inhibition (mm)		
		Reading 1	Reading 2	Average value
<i>Staphylococcus aureus</i>	Positive control <i>Gentamicin</i> (1mg/ml)	9	10	9.5
	Dichloromethane	-	-	-
	Acetone	-	-	-
	Ethanol	-	1	-
	Methanol	2	3	2.5
	Negative control	-	-	-
<i>Streptococcus pyogenes</i>	Positive control <i>Gentamicin</i> (1mg/ml)	15	8	11.5
	Dichloromethane	-	-	-
	Acetone	3	-	-
	Ethanol	3	8	5.5
	Methanol	4	3	3.5
	Negative control	-	-	-

<i>Bacillus cereus</i>	Positive control <i>Gentamicin</i> (1mg/ml)	18	15	16.5
	Dichloromethane	-	-	-
	Acetone	7	4	5.5
	Ethanol	7	2	4.5
	Methanol	4	3	3.5
	Negative control	-	-	-
<i>Escherichia coli</i>	Positive control <i>Gentamicin</i> (1mg/ml)	17	23	20
	Dichloromethane	-	-	-
	Acetone	6	2	4
	Ethanol	7	11	9
	Methanol	5	4	4.5
	Negative control	-	-	-
<i>Acinetobacter baumannii</i>	Positive control <i>Gentamicin</i> (1mg/ml)	16	18	17
	Dichloromethane	-	-	-
	Acetone	7	-	-
	Ethanol	8	2	5
	Methanol	2	2	2
	Negative control	-	-	-
<i>Neisseria gonorrhoeae</i>	Positive control <i>Gentamicin</i> (1mg/ml)	18	16	17
	Dichloromethane	2	-	-
	Acetone	10	2	6
	Ethanol	10	3	6.5
	Methanol	3	2	2.5
	Negative control	-	-	-

#### 4. Discussion

There is an increasing demand for the evaluation of alternative antimicrobial due to the development of resistance to currently available antimicrobial. As per the literature survey, *Cymbopogon citratus* is reported as herbal medicine that is useful in treating various diseases and it is majorly due to its strong antimicrobial activity. Therefore, the investigators planned to work over the In-vitro antimicrobial activity of *Cymbopogon citratus* stem extracts. The antimicrobial study was conducted as per standard reference (Vetrisevan et al., 2012; Fuloria et al., 2020). The antimicrobial results of the present study revealed dichloromethane extracts showed inhibitory effect overgrowth of *Streptococcus pyogenes*, *Bacillus cereus*, *Escherichia coli*, *Acinetobacter baumannii*, *Neisseria gonorrhoeae* in well diffusion method.

Overall, dichloromethane showed more activity against Gram-negative bacteria on well diffusion. Acetone extracts showed inhibitory effect on *Streptococcus pyogenes*, *Bacillus cereus*, *Escherichia coli*, *Acinetobacter baumannii* and *Neisseria gonorrhoeae*. As in disc diffusion, it showed inhibitory effect on *Bacillus cereus*, *Escherichia coli*, *Neisseria gonorrhoeae* except on *A. baumannii*. It showed more antimicrobial activity in the well diffusion method and Gram-negative bacteria. Ethanolic extracts showed an inhibitory effect on *S. pyogenes* and *E. coli* in well diffusion. In disc diffusion, it showed inhibitory effect against *B. cereus*, *E. coli*, *A. baumannii*, and *N. gonorrhoeae*. To summarize, ethanolic extracts showed more antimicrobial activity in the disc diffusion method and Gram-negative bacteria. Methanolic extracts showed an inhibitory effect against *B. cereus* and *N. gonorrhoeae* in well diffusion method. Whereas in disc diffusion method, it showed an inhibitory effect in all studied bacteria. To summarize, methanolic extracts showed more antimicrobial activity in disc diffusion method and Gram-negative bacteria. Based on the present antimicrobial study conducted, the study showed that dichloromethane extract has the highest antimicrobial activity followed by ethanolic extract, acetone extract and lastly methanolic extract. For three microorganisms (*S. pyogenes*, *A. baumannii*, and *N. gonorrhoeae*), so far, no study has been done, it is reported for the first time in our study. Extracts used in the present antimicrobial study have shown overall good antimicrobial activity against these three bacteria in both disc and well diffusion.

## 5. Conclusion

Based on the antibacterial potential of different solvent extracts of *Cymbopogon citratus* stems, it is hereby concluded that dichloromethane extract of *Cymbopogon citratus* stems exerts the highest inhibitory action against the growth of bacterial strains of *Staphylococcus aureus*, *Streptococcus pyogenes*, *Bacillus cereus*, *Escherichia coli*, *Acinetobacter baumannii* and *Neisseria gonorrhoeae*. The high antimicrobial potential of dichloromethane extract of *Cymbopogon citratus* stems may be due to the presence of flavonoidal constituents. The present study concludes the high antibacterial potential of *Cymbopogon citratus* plant. To obtain deeper insight into the mechanism of action of extracts of stems of *Cymbopogon citratus* plant, further investigation on isolates of extracts should be done.

## 6. References

- Ali, B., Al-Wabel, N.A., Shams, S., Ahamad, A., Khan, S.A., and Anwar. F., (2015) Essential oils used in aromatherapy: A systemic review. *Asian Pacific Journal of Tropical Biomedicine*, 5(8), 601-611.
- Azwanida, N.N., (2015) A review on the extraction methods use in medicinal plants, principle, strength and limitation. *Med Aromat Plants*, 4(196), 2167-0412.
- Balouiri, M., Sadiki, M., and Ibnsouda, S.K., (2016) Methods for in vitro evaluating antimicrobial activity: A review. *Journal of Pharmaceutical Analysis*, 6(2), 71-9.
- Calabrese, L., (2005) The use and methods of making a herbarium/plant specimens. A Herb Society of America Guide. *The Herb Society of America*, 6.
- Devillers, J., Steiman, R., and Seigle-Murandi, F., (1989) The usefulness of the agar-well diffusion method for assessing chemical toxicity to bacteria and fungi. *Chemosphere*, 19(10-11), 1693-700.
- Fuloria, S., Ying, C.Y., Xuan, K.Y., Jun, C.T., Karupiah, S., Kumari, U., and Fuloria, N.K. (2020) Determination of total phenolic content and antimicrobial potential of different extracts of *Citrus hystrix* DC leaves, *Bulletin of Environment Pharmacology and life Sciences*, 2020, 9(8), 112-116
- Jaju, S., Patil, A.T, Durgakar, N.J., and Fuloria, N.K. (2009) Phytochemical and antimicrobial activity of stem and leaves of *Desmodium gangeticum* linn", *Hamdard medicus*, 52(4), 131-135.
- Mirghani, M.E., Liyana, Y., and Parveen, J., (2012) Bioactivity analysis of lemongrass (*Cymbopogon citratus*) essential oil. *International Food Research Journal*, 19(2), 569.
- Redfern, J., Kinninmonth, M., Burdass, D., and Verran, J., (2014) Using soxhlet ethanol extraction to produce and test plant material (essential oils) for their antimicrobial properties. *Journal of Microbiology & Biology Education*, 15(1), 45.
- Salome, A.C., Emeka, C.U., Ikechukwu, V.O., Sinye, A.B., Calister, E.U., and Godswill, C.O., (2012) Formulation and evaluation of *Cymbopogon citratus* dried leaf-powder tablets. *African Journal of Pharmacy and Pharmacology*, 6(48), 3274-9.
- Shivkanya, J., Shilpa, P., Sangita, K., and Neeraj, F., (2009) Pharmacognostical studies and antibacterial activity of the leaves of *Murraya koenigii*. *Pharmacognosy Journal*, 1(3), 211-214.
- Srivastava, V., Dubey, S., and Mishra, A. A., (2013) Review on lemon grass: agricultural and medicinal aspect. *International Research Journal of Pharmacy*, 4(8), 42-4.
- Tajidin, N.E., Ahmad, S.H., Rosenani, A.B., Azimah, H., and Munirah, M., (2012) Chemical composition and citral content in lemongrass (*Cymbopogon citratus*) essential oil at three maturity stages. *African Journal of Biotechnology*, 11(11), 2685-93.
- Vetriselvan S, Saminathan, K., Ashok Kumar, J., and Vinoth K., (2019) "Impact of herbal drugs and its clinical application" *International Journal of Research in Pharmaceutical Sciences*, 10(2), 1340-1345.
- Vetriselvan, S., Shankar, J., Gayathiri, S., Ishwin, S., Hemah Devi, C., Yaashini, A., and Sheerenjet, G., (2012) Comparative evaluation of in vitro antibacterial and antioxidant activity using standard drug and polyherbal formulation. *International Journal of Phytopharmacology*, 3(2), 112-116.